



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/34, 43/64, 43/82, A61K 31/165, C07D 207/08		A1	(11) International Publication Number: WO 97/15190
			(43) International Publication Date: 1 May 1997 (01.05.97)
(21) International Application Number: PCT/US96/16865		(US). SHAFER, Jules, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).	
(22) International Filing Date: 21 October 1996 (21.10.96)		(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).	
(30) Priority Data: 60/006,076 24 October 1995 (24.10.95) US 9605163.6 12 March 1996 (12.03.96) GB 60/023,164 5 August 1996 (05.08.96) US		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).		Published With international search report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): LUMMA, William, C. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). TUCKER, Thomas, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). WITHERUP, Keith, M. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). BRADY, Stephen, F. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). WHITTER, Willie, L. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). VACCA, Joseph, P. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). COBURN, Craig [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).			

(54) Title: THROMBIN INHIBITORS

(57) Abstract

A compound which inhibits human thrombin and where has the structure (I) such as (II).



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

- 1 -

TITLE OF THE INVENTION
THROMBIN INHIBITORS

BACKGROUND OF THE INVENTION

5 Thrombin is a serine protease present in blood plasma in the form of a precursor, prothrombin. Thrombin plays a central role in the mechanism of blood coagulation by converting the solution plasma protein, fibrinogen, into insoluble fibrin.

 Edwards *et al.*, *J. Amer. Chem. Soc.* (1992) vol. 114, pp.
10 1854-63, describes peptidyl α -ketobenzoxazoles which are reversible inhibitors of the serine proteases human leukocyte elastase and porcine pancreatic elastase.

 European Publication 363 284 describes analogs of peptidase substrates in which the nitrogen atom of the scissile amide
15 group of the substrate peptide has been replaced by hydrogen or a substituted carbonyl moiety.

 Australian Publication 86245677 also describes peptidase inhibitors having an activated electrophilic ketone moiety such as fluoromethylene ketone or α -keto carboxyl derivatives.

20 Thrombin inhibitors described in prior publications contain sidechains of arginine and lysine. These structures show low selectivity for thrombin over other trypsin-like enzymes. Some of them show toxicity of hypotension and liver toxicity.


 European Publication 601 459 describes sulfonamido
25 heterocyclic thrombin inhibitors, such as N-[4-[(aminoimino-methyl)amino]butyl]-1-[N-(2-naphthalenylsulfonyl)-L-phenylalanyl]-L-prolinamide.

 WO 94/29336 describes compounds which are useful as
30 thrombin inhibitors.

- 2 -

SUMMARY OF THE INVENTION


Compounds of the invention have the following structure:



I

5

and pharmaceutically acceptable salts thereof such as



10

The invention includes a composition for inhibiting loss of blood platelets, inhibiting formation of blood platelet aggregates, inhibiting formation of fibrin, inhibiting thrombus formation, and inhibiting embolus formation in a mammal, comprising a compound of the invention in a pharmaceutically acceptable carrier. These compositions may optionally include anticoagulants (e.g. a fibrinogen receptor antagonist), antiplatelet agents, and thrombolytic agents. The compositions can be added to blood, blood products, or mammalian organs in order to effect the desired inhibitions.

15

The invention also includes a composition for preventing or treating unstable angina, refractory angina, myocardial infarction, transient ischemic attacks, atrial fibrillation, thrombotic stroke, embolic

20

- 3 -

stroke, deep vein thrombosis, disseminated intravascular coagulation, ocular build up of fibrin, and reocclusion or restenosis of recanalized vessels, in a mammal, comprising a compound of the invention in a pharmaceutically acceptable carrier. These compositions may optionally include anticoagulants (e.g. a fibrinogen receptor antagonist), antiplatelet agents, and thrombolytic agents.


The invention also includes a method for reducing the thrombogenicity of a surface in a mammal by attaching to the surface, either covalently or noncovalently, a compound of the invention.

The use of a compound of Claim 1, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for inhibiting thrombus formation, preventing thrombus formation, inhibiting thrombin, inhibiting formation of fibrin, and inhibiting formation of blood platelet aggregates, in a mammal

15

DETAILED DESCRIPTION OF THE INVENTION

Compounds of the invention have the following structure:




I

20

and pharmaceutically acceptable salts thereof wherein

A is



25

- 4 -

wherein

R_a and R_b are independently selected from

hydrogen,

5 a heterocyclic group which is a stable 5- to 7-membered
mono- or bicyclic or stable 7- to 10-membered bicyclic
heterocyclic ring system any ring of which may be
saturated or unsaturated, and which consists of carbon
atoms and from one to three heteroatoms selected from the
10 group consisting of N, O and S, and wherein the nitrogen
and sulfur heteroatoms may optionally be oxidized, and the
nitrogen heteroatom may optionally be quaternized, and
including any bicyclic group in which any of the above-
defined heterocyclic rings is fused to a benzene ring,

15 C_1 -4 alkyl unsubstituted or substituted with CH_3 or C_3 -7
cycloalkyl,

aryl,

substituted aryl with one or two substituents selected from


C_1 -4 alkyl,

20 C_1 -4 alkoxy,
methylenedioxy,
halogen or
hydroxy,

C_3 -7 cycloalkyl,

25 a C_4 -10 carbocyclic or bicyclic ring, or

R_a and R_b , along with the carbon to which they are attached,
form a C_3 -7 cycloalkyl ring or




30 where R^{10} is H or -OH, and
 R^{11} is H or -OCH₃, and

X is -NHR_c or -OH, wherein,

- 5 -


 R_C is

hydrogen,

-CH₃,-(CH₂)₁₋₃CH₃,5 -(CH₂)₂₋₄OH,-(CH₂)₁₋₃COOH,-(CH₂)₁₋₃COOR⁶, where R⁶ is C₁₋₄alkyl,-(CH₂)₁₋₃CONR⁷R⁸,10 where R⁷ and R⁸ are independently hydrogen or C₁₋₄alkyl,

where D is 1, 2, 3, or 4 carbon atoms unsubstituted

or any 1, 2, 3, or 4 of which are substituted with OH,


-SO₂(CH₂)₁₋₃aryl,15 -(CH₂)₁₋₃NH₂,C₃₋₇ cycloalkyl ring unsubstituted or substituted with-OH, -C(O)OH, or -C(O)OR_d, where R_d isC₁₋₄ alkyl,

20





Y is O or NH,

W is C or N,


Z is C or N, and

25 R⁶ is -CH₂OH or -N(CH₃)₂ provided that W and Z are not the same,R⁷ is H or CH₃, and

- 6 -

R⁸ is H or


5

where R⁹ is H, NH₂, or OH;

or

10

A is



, wherein

15

B is a bond, O, -CH₂-O-, or -O-CH₂-;


R² and R⁵ are independently selected from
hydrogen, provided that R² and R⁵ are not both hydrogen,
C₁₋₄ alkyl,
20 C₁₋₄ alkoxy,
halogen,

- 7 -

- COOH,
 -OH,
 -COOR⁶, where R⁶ is C₁-4alkyl,
 -CONR⁷R⁸, where R⁷ and R⁸ are independently
 5 hydrogen or C₁-4alkyl,
 -OCH₂CO₂H,
 -OCH₂CO₂CH₃,
 -OCH₂CO₂(CH₂)₁₋₃CH₃,
 -O(CH₂)₁₋₃C(O)NR³R⁴, wherein R³ and R⁴ are independently
 10 hydrogen, C₁-4alkyl, C₃-7 cycloalkyl, or -CH₂CF₃,
 -(CH₂)₁₋₄OH,
 -NHC(O)CH₃,
 -NHC(O)CF₃,
 -NHSO₂CH₃, and
 15 -SO₂NH₂; and

m is 1 or 2.

In one class, the compounds have the following structure:



20

and pharmaceutically acceptable salts thereof wherein

X is as previously defined,

25


R_a and R_b are as previously defined,

R² and R⁵ are as previously defined, and

- 8 -

m is as previously defined.

A first subclass of this class of compounds has the formula




and pharmaceutically acceptable salts thereof, wherein



R² is -OCH₂C(O)NHR⁴; and

10 R⁴ is -CH₂CH₃, cyclopropyl, or -CH₂CF₃.

Examples of compounds in the first subclass include




- 9 -



5 and pharmaceutically acceptable salts thereof.

A second subclass of this class of compounds has the formula



10

and pharmaceutically acceptable salts thereof wherein

X is -NHR_C or -OH, wherein

- 10 -

 R_c is

hydrogen,

-CH₃,-(CH₂)₁₋₃CH₃,5 -(CH₂)₂₋₄OH,-(CH₂)₁₋₃COOH,-(CH₂)₁₋₃COOR⁶, where R⁶ is C₁₋₄alkyl,-(CH₂)₁₋₃CONR⁷R⁸, where R⁷ and R⁸ are independently hydrogen or C₁₋₄alkyl,10 $-(CH_2)_{1-3}CON \begin{array}{c} \diagup \diagdown \\ \text{D} \end{array}$,

where D is 1, 2, 3, or 4 carbon atoms unsubstituted or any 1, 2, 3, or 4 of which are substituted with OH,

-SO₂(CH₂)₁₋₃aryl,-(CH₂)₁₋₃NH₂,15 C₃₋₇ cycloalkyl ring unsubstituted or substituted with -OH,-C(O)OH, or -C(O)OR_d, where R_d is C₁₋₄ alkyl,
$$-(CH_2)_{1-3} \begin{array}{c} \text{Y} \\ \diagup \quad \diagdown \\ \text{W} \quad \text{Z} \end{array} \text{R}^6$$

where

20 Y is O or NH,

W is C or N,






Z is C or N, and

R⁶ is -CH₂OH or -N(CH₃)₂ provided that W and Z are not the same,
$$25 \quad -(CH_2)_{1-3} \begin{array}{c} \text{O} \\ \parallel \\ \text{C} \end{array} \text{N} \begin{array}{c} \text{R}^8 \\ \diagup \quad \diagdown \\ \text{NR}^7 \end{array}$$

where

R⁷ is H or CH₃, andR⁸ is H or

- 11 -



where R^9 is H, NH_2 , or OH;


R_a and R_b are as previously defined, and

10

R^2 and R^5 are independently selected from
hydrogen, provided that R^2 and R^5 are not both hydrogen,
C₁₋₄ alkyl,
C₁₋₄ alkoxy,
15 halogen, and
—OH.

A group of this second subclass of compounds has the
formula

20



- 12 -


and pharmaceutically acceptable salts thereof wherein

X is as previously defined,

- 5 R_a and R_b are independently selected from
hydrogen,
a heterocyclic group which is a stable 5- to 7-membered mono- or
bicyclic or stable 7- to 10-membered bicyclic heterocyclic ring
system any ring of which may be saturated or unsaturated, and
10 which consists of carbon atoms and from one to three
heteroatoms selected from the group consisting of N, O and S,
and wherein the nitrogen and sulfur heteroatoms may
optionally be oxidized, and the nitrogen heteroatom may
optionally be quaternized, and including any bicyclic group in
15 which any of the above-defined heterocyclic rings is fused to a
benzene ring,
C₁₋₄ alkyl unsubstituted or substituted with CH₃ or C₃₋₇
cycloalkyl,
phenyl, or
20 R_a and R_b , along with the carbon to which they are attached,
form a cyclohexyl ring; and
 R^2 and R^5 are independently selected from
hydrogen, provided that R^2 and R^5 are not both hydrogen,
Cl,
25 -CH₃,
-CH₂CH₃,
-OCH₃, and
-OH.

- 30 One subgroup of this group of compounds has the formula

- 13 -




and pharmaceutically acceptable salts thereof wherein


5 R² and R⁵ are independently selected from -OCH₃ and -CH₃; and

R_C is hydrogen or -SO₂CH₂C₆H₅.


Examples of this subgroup include



10




- 14 -



and pharmaceutically acceptable salts thereof.

5 A second subgroup of this group of compounds has the formula




10 and pharmaceutically acceptable salts thereof wherein

X is as previously defined, and

R_a and R_b are as previously defined.

15 A family of the second subgroup of compounds has the formula

- 15 -




and pharmaceutically acceptable salts thereof, wherein

5 R_c is

hydrogen,

$SO_2CH_2C_6H_5$, or

10






; and

R_a and R_b are phenyl, or R_a and R_b , along with the carbon to which they are attached, form cyclohexyl.



15

Examples of the family include

- 16 -



- 17 -



and pharmaceutically acceptable salts thereof.

5

Some abbreviations that may appear in this application are as follows.

<u>Designation</u>	
10 BOC (Boc)	t-butyloxycarbonyl
HBT(HOBT or HOBT)	1-hydroxybenzotriazole hydrate
BBC reagent	benzotriazolyloxy-bis(pyrrolidino)- carbonium hexafluorophosphate
PyCIU	1,1,3,3-bis(tetramethylene)- chlorouronium hexafluorophosphate
15 EDC	1-ethyl-3-(3-dimethylaminopropyl)
	carbodiimide hydrochloride
(BOC) ₂ O	di-t-butyl dicarbonate
DMF	dimethylformamide
20 Et ₃ N or TEA	triethylamine

- 18 -

	EtOAc	ethyl acetate
	TFA	trifluoroacetic acid
	DMAP	dimethylaminopyridine
	DME	dimethoxyethane
5	BH ₃ -THF	Borane-tetrahydrofuran complex
	D-Phe(3,4-Cl ₂)	D-3,4-Dichlorophenylalanine
	D-3,3-dicha	D-3,3-Dicyclohexylalanine
	Pro	Proline
	Arg	Arginine
10	Gly	Glycine
	D-3,3,-diphe	D-3,3-Diphenylalanine

The compounds of the present invention may have chiral centers and occur as racemates, racemic mixtures and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention.

When any variable occurs more than one time in any constituent or in formula I, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

The term "aryl" means a 5- or 6-membered aromatic ring containing 0, 1, or 2 heteroatoms selected from O, N, and S. Examples of aryl include phenyl, pyridine, pyrimidine, imidazole, thiophene, oxazole, isoxazole, thiazole, and amino- and halogen- substituted derivatives thereof.

The term "alkyl" means straight or branched alkane containing 1 to about 10 carbon atoms, e.g., methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl, octyl radicals and the like, straight or branched alkene containing 2 to about 10 carbon atoms, e.g., propylenyl, buten-1-yl, isobutenyl, pentenylen-1-yl, 2,2-methylbuten-1-yl, 3-methylbuten-1-yl, hexen-1-yl, hepten-1-yl, and octen-1-yl radicals and the like, or straight or branched alkyne containing 2 to about 10 carbon atoms, e.g., ethynyl, propynyl,

- 19 -

butyn-1-yl, butyn-2-yl, pentyn-1-yl, pentyn-2-yl, 3-methylbutyn-1-yl, hexyn-1-yl, hexyn-2-yl, hexyn-3-yl, 3,3-dimethylbutyn-1-yl radicals and the like.

5 The term "alkoxy" represents an alkyl group of indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include methyloxy, propyloxy, and butyloxy.

The terms "Halo" or "halogen," as used herein, means fluoro, chloro, bromo and iodo.

10 The term "counterion" is used to represent a small, single negatively-charged species, such as chloride, bromide, hydroxide, acetate, trifluoroacetate, perchlorate, nitrate, benzoate, maleate, tartrate, hemitartrate, benzene sulfonate, and the like.


15 The term heterocycle or heterocyclic, as used herein except where noted, represents a stable 5- to 7-membered mono- or bicyclic or stable 7- to 10-membered bicyclic heterocyclic ring system any ring of which may be saturated or unsaturated, and which consists of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom
20 may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure.

Examples of such heterocyclic elements include piperidinyl, piperazinyl,
25 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodiny, 2-oxoazepinyl, azepinyl, pyrrolyl, 4-piperidonyl, pyrrolidinyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl,
30 quinuclidinyl, isothiazolidinyl, indolyl, quinolinyl, isoquinolinyl, benzimidazolyl, thiadiazoyl, benzopyranyl, benzothiazolyl, benzoxazolyl, furyl, tetrahydrofuryl, tetrahydropyranyl, thienyl, benzothienyl, thiamorpholinyl, thiamorpholinyl sulfoxide,

- 20 -

thiamorpholinyl sulfone, and oxadiazolyl. Morpholino is the same as morpholinyl.

An example of the moiety of R_a or R_b independently selected from substituted aryl with one or two substituents selected from methylenedioxy is



The pharmaceutically-acceptable salts of the compounds of Formula I (in the form of water- or oil-soluble or dispersible products) include the conventional non-toxic salts or the quaternary ammonium salts which are formed, e.g., from inorganic or organic acids or bases. Examples of such acid addition salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, and undecanoate. Base salts include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups may be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides like benzyl and phenethyl bromides and others.

Amide couplings used to form the compounds of this invention are typically performed by the carbodiimide method with


- 21 -

reagents such as dicyclohexylcarbodiimide, or 1-ethyl-3-(3-dimethyl-aminopropyl) carbodiimide. Other methods of forming the amide or peptide bond include, but are not limited to the synthetic routes via an acid chloride, azide, mixed anhydride or activated ester. Typically,
5 solution phase amide coupling are performed, but solid-phase synthesis by classical Merrifield techniques may be employed instead. The addition and removal of one or more protecting groups is also typical practice.

Compounds of the invention can be prepared according to
10 the general procedures outlined below:


A protected amino acid such as D-cyclohexylglycine is coupled to proline methyl ester using a coupling agent such as EDC and HOBT. The coupled product is then hydrolyzed with base such as lithium hydroxide, and the resultant acid is coupled to the desired amine
15 such as 2,5-dichlorobenzylamine. The product is treated with a strong acid such as HCl gas or trifluoroacetic acid to remove the t-butyloxycarbonyl protecting group. Tables I and II illustrate compounds synthesized in this manner and are exemplified by Example 1.

- 22 -

SCHEME 1

- 5 A method for synthesizing compounds illustrated in tables 2 and 3 is to react a free amino containing compound with an alkylating agent such as t-butyl-bromoacetate. The resulting compound is treated with acid to form an acid, and the resultant acid is coupled to the desired amine under standard coupling conditions. If the product has a protecting group, this is conveniently removed with acid (for acid lable groups).
- 10


- 23 -

SCHEME 2

- 24 -


An alternate method for functionalizing the amine group is illustrated in Scheme 3. An amine, such as that from Example 1, treated with an aldehyde and a reducing agent such as sodium triacetoxyborohydride to give the desired product.

5

SCHEME 3


- 10 β -Aminoalkylsulfonamide containing compounds are synthesized by
reacting an amino compound with a sulfonylating reagent such as
chloroethyl sulfonyl chloride and a base such as triethylamine. The
product is reacted with a primary or secondary amine to give the
product. In some cases the amine contains a protecting group which is
15 removed with acid.

- 25 -


SCHEME 4

- 26 -

SCHEME 5



- 27 -



5

R represents, for example, hydrogen, C₁-4 alkyl, C₁-4 cycloalkyl or CH₂CF₃.

- 28 -

EXAMPLE 1Preparation of D- β , β -diphenylala-Pro-N-(2,5-dichlorophenyl)methyl amide (1-1)

- 5 A solution of 418.00 mg (0.95 mmol) of Boc-(D)- β , β -diphenylala-ProOH, 168.00 mg (0.95 mmol) of 2,5-dichlorobenzylamine, 201.00 mg (1.05 mmol) of EDC, 142.00 mg (1.05 mmol) of HOBT, and 146.00 ml (1.05 mmol) of triethylamine in 8 ml anh. DMF was stirred at room temp. in an argon atmosphere for 18 h. The
10 reaction was diluted with three times its volume of aq. 10% citric acid solution, and the resulting suspension was stirred vigorously for 45 min. The suspension was filtered, and the solid product dried *in vacuo* over anh. P₂O₅ to give 540 mg of the intermediate coupling product. The solid was dissolved in a minimum of EtOAc with a small amt. of CHCl₃
15 added to aid dissolution. The solution was cooled to -10°C, and was bubbled with HCl gas for approx. five minutes. The solution was stirred at this temp. for twenty additional minutes, and was removed from the cooling bath. The solution was purged with argon, and a white amorph. solid precipitate resulted. Filtration and drying provided 1-1
20 as a white powder. Anal.(C₂₇H₂₇N₃O₂Cl₂ • HCl • 0.35 H₂O • 0.50 CHCl₃), CHN. High res. MS: theo., 496.15585; obs., 496.15652.

EXAMPLE 2

- 25 Preparation of D- β , β -diphenylala-Pro-N-(2-hydroxy-5-methyl)-benzylamide (2-1)

- A solution of 96 mg (0.22 mmol) of Boc-D-diphenylala-Pro-OH and 40 mg (0.20 mmol) of 2-hydroxy-5-ethyl benzylamine hydrochloride in 15 ml of DMF was treated with 37 mg (0.24 mmol) of
30 HOBT-H₂O and N-methyl morpholine (pH 8 moistened pH 5-10 paper) followed by 50 mg of EDC (0.26 mmol). After stirring overnight, the reaction mixture was evaporated to dryness, the residue partitioned with EtOAc/dilute NaHCO₃; the organic layer washed with H₂O, dilute NaHCO₃, sat'd. NaCl; solvent was removed to afford crude

- 29 -

intermediate. Approx. 3 ml of 100% trifluoroacetic acid was added to the residue, the solution set 15 min; the TFA was evaporated *in vacuo* and replaced with CH₃CN-CH₃OH-H₂O (1:1:3), followed by preparative HPLC to afford, after lyophilization of fractions, 2-1.

5 FAB-MS m/c 472 (M+H); HPLC >99%.

EXAMPLE 3

Preparation of D-β,β-diphenylala-Pro-N-(2,5-dimethoxy)-benzylamide
10 (3-1)

A solution of 242 mg (0.55 mmol) of BOC-D-diphenylala-Pro-OH and 84 mg (0.50 mmol) of 2,5-dimethoxy benzylamine in 20 ml of DMF was treated with 92 mg (0.80 mmol) of HOBT, N-methylmorpholine, and 125 mg (0.65 mmol) of EDC as in Example 2.

15 Standard workup afforded crude intermediate which treated with 5 ml of 100% TFA to remove the BOC group as in Example 2. Preparative HPLC afforded 170 mg of the desired product as the TFA salt, which was converted to the HCl salt. (HCl/EtOAc) to afford 3-1: FAB-MS m/e 488 (M+H), HPLC ca. 90%.

20

EXAMPLE 4

Preparation of N-carboxymethyl-D-β,β-diphenylala-Pro-N-(2,5-dimethoxy)-benzylamide (4-1)

25 A solution of 40 mg (0.082 mmol) of 3-1 and 16 mg of t-butyl bromoacetate with 22 ml (1.5 equiv.) of DIEA in 0.5 ml of DMF was stirred 20 min at 25°; followed by an additional equal amount of the latter two reagents, the reaction was complete in 48 hrs. After dilution with EtOAc, extractive workup afforded 38 mg of glassy solid
30 intermediate. Approx. 3 ml of 100% TFA was used to remove the t-butyl ester, as in Example 2; the compound was purified by semi-preparative HPLC and the pooled fractions were evaporated and converted to the HCl salt. Filtration of the precipitated HCl salt from hexane•EtOAc gave 4-1. FAB-MS m/e 546 (M+H), HPLC ca. 95%.

- 30 -

EXAMPLE 5Preparation of N-[2-(imidazolyl)-methyl]-D- β , β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide (5-1)

5 A solution of 107 mg (0.20 mmol) of 1-1 in 2.0 ml of 0.24 M HOAc in 1,2-dichloroethane under N₂ was treated with 21 mg (0.21 mmol) of imidazole-2-carboxaldehyde, followed by 64 mg (0.30 mmol) of sodium triacetoxyborohydride. After 4 days an additional 0.5 equivalents more of the latter reagents were added, and the reaction was
10 stirred an additional 2 days. The mixture was concentrated *in vacuo* to dryness, dissolved in ca. 1;3 HOAc-H₂O, and purified by preparative HPLC. Pooling of product containing fractions yielded, after lyophilization, 5-1: FAB-MS m/e 576 (M+H); HPLC ca. 95%.

15

EXAMPLE 6Preparation of N-[4-(imidazolyl)-methyl]-D- β , β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide (6-1)

20 As in Example 5 above, a solution of 214 mg (0.40 mmol) of 1-1 in 4.0 ml of 1,2-dichloroethane was treated with 59 mg (0.60 mmol) of imidazole-4-carboxaldehyde and 176 mg (0.80 mmol) of sodium triacetoxyborohydride. After 24 h. the solvent was concentrated *in vacuo* and the product purified by preparative HPLC as above to yield 141 mg of lyophilized 6-1: FAB-MS m/e 576 (M+H); HPLC 99%.

25

EXAMPLE 7Preparation of N-[2-(5-hydroxymethylfuryl)-methyl]-D- β , β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide (7-1)

30 As in Example 5 above, a solution of 204 mg (0.40 mmol) of 1-1 in 4.0 ml of 1,2-dichloroethane under N₂ was treated with 74 mg (0.60 mmol) of 5-hydroxymethyl-2-furaldehyde and 164 mg (0.80 mmol) with sodium triacetoxyborohydride. After 24 hr the solvent was

- 31 -

concentrated *in vacuo* and the product purified by preparative HPLC as above to yield 7-1: FAB-MS m/e 606 (M+H); HPLC 99%.

EXAMPLE 8

5

Preparation of N-[2-(5-dimethylaminofuryl)-methyl]-D- β,β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide (8-1)

As in Example 5 above, a solution of 206 mg (0.40 mmol) of 1-1 in 4.0 ml of 1,2-dichloroethane under N₂ was treated with 86 mg (0.60 mmol) of 5-dimethylamino-2-furaldehyde and 170 mg (0.80 mmol) of sodium triacetoxyborohydride. After 24 hr the solvent was concentrated *in vacuo* and the product purified by preparative HPLC as above to yield 8-1: FAB-MS m/e 619 (M+H); HPLC >99%.

15

EXAMPLE 9

Preparation of N-(imino-aminomethyl)-methyl-D- β,β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide (9-1)

A solution of 20 mg of 1-1 in 2.0 ml of DMF was treated with 11 mg of chloroacetamide hydrochloride, followed by 2 drops of diisopropyl ethylamine. The mixture was heated at 50-60° for 2 days; the solvent was evaporated *in vacuo* and the residue in H₂O/5% acetonitrile was processed by preparative HPLC to yield, after lyophilization, 9-1: FAB-MS m/e 552 (M+H); HPLC ca. 88%.

25

EXAMPLE 10

Preparation of D-cyclohexylglycyl-Pro-N-(2,5-dichloro)-benzylamide (10-1)

A solution of 1.00 g (3.89 mmol) of Boc-D-cyclohexyl glycine and 1.26 g (4.08 mmol) of (H)-prolyl-2,5-dichlorobenzylamide hydrochloride in 90 ml of DMF was treated with 0.71 g (4.67 mmol) of HOBt·H₂O and N-methyl morpholine (pH 8); then 0.97 g (5.06 mmol) of EDC, followed by stirring 5 hr. The solution was concentrated *in*

- 32 -

vacuo to a volume of ca. 20 ml, followed by partition with EtOAc/dilute NaHCO₃ and extractive workup as in Example 2 to give crude intermediate, which was purified by chromatography on silica gel, eluting with 1:1 EtOAc/hexane, to give 1.87 g (94% yield of coupled intermediate). The above sample in approx. 50 ml of EtOAc was saturated with HCl gas at -10°, set 60 min at 0-20°, followed by purging with N₂, as precipitate slowly formed. The solid was filtered and washed with ether, drying *in vacuo* to give 10-1: FAB-MS m/3 413 (M+H); HPLC 97%.

10

EXAMPLE 11

Preparation of N-carboxymethyl-D-cyclohexylglycyl-Pro-N-(2,5-dichloro)-benzylamide (11-1)

15 A solution of 289 mg (0.70 mmol) of 10-1 and 0.23 ml (0.28 g, 1.44 mmol) of *t*-butyl bromoacetate with 0.24 ml of DIEA in 5.0 ml of DMF, was stirred at 25° for 2 days. The solvent was removed *in vacuo*, the residue partitioned with EtOAc/dilute NaHCO₃, and the organic layer was washed with saturated NaCl and dried over Na₂SO₄.
20 Solvent removal afforded 390 mg of crude intermediate, HPLC 95%. A solution of 87 mg of the above intermediate in 10 ml of EtOAc/CH₂Cl₂ (4/1) was saturated with HCl at -10°, set 30 min; then purged with N₂, and the solution concentrated under reduced pressure until appearance of solid. Precipitation was completed by addition of ether, and product
25 was isolated by filtration, washing with ether, and drying *in vacuo* to give 11-1: FAB-MS m/e 480 (M+H); HPLC ca. 90%.

EXAMPLE 12

30 Preparation of N-((1-piperazinyl)-carboxy)-methyl-D-cyclohexylglycyl-Pro-N-(2,5-dichloro)-benzylamide (12-1)

A solution of 80 mg (0.16 mmol) of 11-1 and 36 mg (0.19 mmol) of *t*-BOC-1,4-piperazine in 2.0 ml of DMF was treated with 32 mg (0.21 mmol) of HOBt•H₂O and N-methyl morpholine (pH 8); then

- 33 -

43 mg (0.22 mmol) of EDC was added, followed by stirring at 25° for 20 hr. The solvent was evaporated *in vacuo* and the residue partitioned with EtOAc/dilute NaHCO₃, washing with 2 portions of saturated NaCl, and dried over Na₂SO₄. Removal of solvent afforded 100 mg of crude
5 intermediate. To the above sample was added 5.0 ml of TFA; let stir for 30 min, the TFA was evaporated under reduced pressure and the product was purified to give lyophilized 12-1; FAB-MS m/e 538 (M+H); HPLC 97%.

10

EXAMPLE 13

Preparation of D-β,β-diphenylala-Pro-N-(2,5-dimethylbenzyl)amide
(13-1)

In a similar manner as in Example 1 but substituting 2,5-
15 dimethylbenzylamine for 2,5-dichlorobenzylamine, 13-1.

EXAMPLE 14

Preparation of N-Phenylmethanesulfonyl-D-β,β-diphenylala-Pro-N-
20 (2,5-dimethylbenzyl)amide (14-1)

D-β,β-diphenylala-L-Pro-N-(2,5-dimethylbenzyl)amide
hydrochloride is reacted with hexamethyldisilazane (0.10 ml per 32 mg
hydrochloride) in dry acetonitrile for 5 min at reflux. The mixture is
cooled 30 min at room temperature and treated with phenylmethane-
25 sulfonyl chloride (50 mg). After 15 min at room temperature the
mixture is diluted with CH₂Cl₂. The CH₂Cl₂ solution is washed with
water, dried (Na₂SO₄) filtered and concentrated *in vacuo*.

Chromatography on activity III neutral alumina gave 14-1. M⁺H⁺/e 610
(calc'd for (C₃₆H₃₉N₃O₄S) = 609.794.

30

- 34 -

EXAMPLE 15Preparation of N-(4-pyridylmethanesulfonyl)-D-β,β-diphenylala-Pro-N-(2,5-dichlorobenzyl)amide (15-1)

- 5 In a similar manner 1-1 is reacted with 43 mg 4-pyridylmethanesulfonyl chloride (trifluoromethanesulfonic acid salt) and hexamethyldisilazane. Similar workup and preparative HPLC gave lyophilized fractions of the title compound as the trifluoroacetic acid salt. This is treated with HCl(g) in ethyl acetate to give the crystalline
10 hydrochloride of 15-1; high resolution MS (M^+H^+/e) = 651.605 ($C_{33}H_{32}Cl_2N_4O_4S^+H^+$).

EXAMPLE 16

- 15 Preparation of N-[(N,N-diethylcarboxamido)methyl]-D-β,β-diphenylala-Pro-N-(2,5-dichloro)-benzylamide (16-1)

- A solution of 100.00 mg (0.19 mmol) of 1-1, 41.00 mg (0.21 mmol) of alpha-bromo-(N,N-diethyl)acetamide, and 75.00 ml (0.42 mmol) of diisopropylethylamine in 1 ml anh. DMF was stirred at 50°C
20 in an argon atmosphere for 4 h. The solution was further stirred at room temp. for 48 h, and was concentrated *in vacuo* to give a tan oil. The crude oil was purified via reverse phase prep LC, and the pure product fractions combined and lyophilized. Lyophilization provided 16-1 as a fluffy white amorphous solid. Anal. ($C_{33}H_{38}N_4O_3Cl_2 \cdot 2.00$
25 TFA $\cdot 1.00 H_2O$), CHN. Mass Spec.: $M^+ = 609$.

EXAMPLE 17

- 30 Preparation of N-[(4-methylpiperazine)carboxamidomethyl]-D-β,β-diphenylala-Pro-N-(2,5-dichloro)-benzylamide (17-1)

A solution of 38.00 mg (0.06 mmol) of N-carboxymethyl-D-β,β-diphenylala-Pro-N-(2,5-dichloro)-benzylamide, 7.00 ml (0.06 mmol) of 4-methyl piperazine, 1.00 mg (1.10 mmol) of EDC, 10.00 mg (1.10 mmol) of HOBT, and 20.00 ml (2.20 mmol) of triethylamine in 1

- 35 -

ml of anh. DMF was stirred for 18 h in an argon atm. The reaction was concentrated *in vacuo* to give a clear oil, which was purified via reverse phase prep LC. Pure product fractions were combined and lyophilized to provide 17-1 as an amorphous white powder.

- 5 Anal.(C₃₄H₃₉N₅O₃Cl₂ • 2.15 TFA • 2.20 H₂O), CHN. Mass Spec.:
M⁺ = 636.

EXAMPLE 18

- 10 Preparation of D-β,β-diphenylala-Pro-N-(2-hydroxy-5-chloro)-
benzylamide (18-1)

- A solution of 278.00 mg (0.64 mmol) of Boc-D-β,β-diphenylala-ProOH, 100.00 mg (0.64 mmol) of 2-hydroxy-5-chlorobenzylamine, 136.00 mg (0.71 mmol) of EDC, 96.00 mg (0.71
15 mmol) of HOBT, and 99.00 ml (0.71 mmol) of triethylamine in 2 ml anh. DMF was stirred in an argon atm. for 18 h. The reaction was diluted with aq. 10% citric acid, and the resulting suspension was stirred vigorously for 45 min. The suspension was filtered, and the recovered white solid dried *in vacuo*. The solid was dissolved in a minimum of
20 EtOAc, and the solution was cooled to -10°C. The solution was bubbled with HCl gas for approx. five minutes, and was stirred for an additional 30 min. The reaction was removed from the cooling bath, and was purged with argon. The solution was concentrated *in vacuo* to provide a clear oil. The oil was purified via reverse phase prep LC, and the pure
25 product fractions combined and lyophilized to give 18-1 as a fluffy white amorphous powder. Anal. (C₂₇H₂₈N₃O₃Cl • 1.30 TFA • 0.55 H₂O), CHN. High Res. MS: theo. = 478.18975, obs. = 478.18940.

EXAMPLE 19

- 30 Preparation of N-[(N,N-diethylcarboxamido)methyl]-D-β,β-diphenylala-Pro-N-(3-chloro)-benzylamide (19-1)

 A solution of 150.00 mg (0.30 mmol) of D-β,β-diphenylala-Pro-N-(3-chloro)-benzylamide HCl (prepared from Boc-

- 36 -

(D)-Dip-ProOH and 3-chlorobenzylamine via a procedure analogous to that described in Example 1), 64.00 mg (0.33 mmol) of alpha-bromo-N,N-diethylacetamide, and 105.00 ml (0.60 mmol) of triethylamine in 1 ml of anh. DMF was stirred at room temp. in an argon atm. for 18 h.

- 5 The reaction was concentrated *in vacuo* to give a brown oil. The crude oil was purified by reverse phase prep LC, and the pure product fractions combined and lyophilized to give 19-1 as a tacky white amorphous powder. Anal. (C₃₃H₃₉N₄O₃Cl • 1.65 TFA • 0.10 H₂O), CHN. Mass Spec.: M⁺ = 575.

10

EXAMPLE 20

Preparation of α -(R)-amino- α -(3,4-methylenedioxybenzyl)acetyl-Pro-N-(2,5-dichloro)-benzylamide (20-1)

- 15 A solution of 100.00 mg (90.30 mmol) of α -(R)-azido- α -(3,4-methylenedioxybenzyl)acetyl-ProOH, 53.00 mg (0.30 mmol) of 2,5-dichlorobenzylamine, 63.00 mg (0.33 mmol) of EDC, 45.00 mg (0.33 mmol) of HOBT, and 47.00 ml (0.33 mmol) of triethylamine in 1 ml of anh. DMF was stirred at room temp. in an argon atm. for 18
- 20 h. The reaction was diluted with 3 times its volume of aq. 10% citric acid, and the solution stirred for approx. 10 min. The mixture was extracted with 2 x 25 ml of EtOAc, and the combined extracts washed with water and brine and dried over anh. MgSO₄. Concentration provided a foam, which was purified via gravity column
- 25 chromatography over silica gel with 2.5% MeOH/CHCl₃. Concentration of the pure fractions provided 120 mg of coupling product as a white foam. The coupling product (120.00 mg/0.27 mmol) was dissolved in 3 ml of THF to which was added 50 ml of H₂O. The solution was treated with 71.00 mg (0.27 mmol) of triphenylphosphine,
- 30 and the resulting solution stirred at 55°C for 18 h. The reaction was concentrated *in vacuo* to a clear oil, which was purified via reverse phase prep LC. The pure product fractions were combined and lyophilized to provide 20-1 as a tacky white amorphous powder. Anal.

- 37 -

(C₂₂H₂₁N₃O₄Cl₂ • 1.05 TFA • 1.00 H₂O), CHN. Mass spec.: M⁺ = 464.

EXAMPLE 21

5

Preparation of D,L-(3,4-methylenedioxy)phenylglycine-Pro-N-(2,5-dichloro)-benzylamide (21-1)

10 A solution of 100.00 mg (0.34 mmol) of Boc-D,L-(3,4-methylenedioxy)phenylglycine, 105.00 mg (0.34 mmol) of 2,5-dichlorobenzylamine, 73.00 mg (0.38 mmol) of EDC, 51.00 mg (0.38 mmol) of HOBt, and 105.00 ml (0.75 mmol) of triethylamine in 2 ml of anh. DMF was stirred for 18 h in an argon atmosphere. The reaction was diluted with 4 times its volume of aq. 10% citric acid, and the resulting suspension stirred vigorously for approx. 45 min. The

15 suspension was filtered to give a white solid which was dried *in vacuo* over P₂O₅ to give 185 mg of crude coupling product. The product from above was dissolved in a min. of EtOAc, and the solution cooled to -10°C. The cold solution was bubbled with HCl gas for approx. 5 min., and was stirred in the cold for an additional 20 min. The reaction was

20 removed from the bath, and was purged with argon. A white precip. resulted, which was isolated via filtration. The solid became extremely tacky on exposure to the air, and was redissolved in EtOAc and dried over anh. MgSO₄. The solution was concentrated to an off-white oil/solid. The crude product was purified via reverse phase prep LC,

25 and the pure product fractions combined and lyophilized. Lyophilization provided 21-1 as a fluffy white amorphous powder which was determined by HPLC to be a 1:1 mixture of diastereomers at the phenylglycine center. Anal.(C₂₁H₂₁N₃O₄Cl₂ • 1.30 TFA • 0.10 H₂O), CHN. Mass Spec.: M⁺ = 450.

30

- 38 -

EXAMPLE 22

Preparation of N-[(N,N-diethylcarboxamido)methyl]-(D)-cyclohexylglycine-Pro-N-(2,5-dichloro)-benzylamide (22-1)

- 5 A solution of 50.00 mg (0.11 mmol) of D-cyclohexylglycine-Pro-N-(2,5-dichloro)-benzylamide HCl, 21.40 mg (0.11 mmol) of alpha-bromo-N,N-diethylacetamide, and 38.20 ml (0.22 mmol) of diisopropylethylamine in 1 ml anh. DMF was stirred in an argon atm. for 18 h. HPLC indicated that the reaction was only approx. 50%
10 complete, so an additional 0.50 equivalents of the bromide was added, and the reaction was warmed to 60°C for approx. 4 h. The reaction was concentrated *in vacuo*, and the crude brown oil product purified via reverse phase prep LC. Pure product fractions were combined and lyophilized to provide 22-1 as a tacky white amorphous powder. Anal.
15 (C₂₆H₃₈N₄O₃Cl₂ • 1.65 TFA • 0.65 H₂O), CHN. Mass Spec.: M⁺ = 525.

EXAMPLE 23

- 20 Preparation of D-cyclohexylglycine-homopro-N-(2,5-dichloro)-benzylamide (23-1)

- A solution of 199.00 mg (0.77 mmol) of Boc-D-cyclohexylglycine, 250.00 mg (0.77 mmol) of proline-N-(2,5-dichloro)-benzylamide, 163.00 mg (0.85 mmol) of EDC, 115.00 mg (0.85 mmol)
25 of HOBT, and 237.00 ml (1.70 mmol) of triethylamine in 5 ml of anh. DMF was stirred for 18 h in an argon atmosphere. The reaction was diluted with 3 times its volume of aq. 10% citric acid, and the resulting suspension stirred vigorously at room temp. for approx. 90 min. The suspension was filtered and the white solid dried *in vacuo* to provide
30 321 mg of the crude coupling product. The coupling product was dissolved in a min. of EtOAc, with a small amt. of CHCl₃ added to assist in solubilizing the material. The reaction was cooled to -10°C, and was bubbled with HCl gas for approx. 10 min. The cold solution was stirred for an additional 30 min., and the bath removed. The reaction was

- 39 -

purged with argon, which provided a precipitate. Filtration and drying *in vacuo* provided 23-1 as a white crystalline solid, MP = 198-201°C. Anal. (C₂₁H₂₉N₃O₂Cl₂ • HCL • 1.05 H₂O • 1.25 CHCl₃), CHN. Mass spec.: M⁺ = 426.

5

EXAMPLE 24

Preparation of N-(2-(1-pyrrolidinyl)-ethanesulfonyl)-amino-D-β,β-diphenylala-Pro-N-(2,5-dichlorobenzyl)amide (24-1)

10 A cooled suspension of 250.00 mg (0.43 mmol) 1-1 in dichloromethane is treated with three equivalents of triethylamine. The reaction is allowed to warm to room temperature over 18 hrs and is then concentrated and chromatographed via preparative TLC. The product is dissolved in acetonitrile which contains 2 equivalents of
15 pyrrolidine. After stirring at room temperature for 48 hr, the reaction is concentrated and 24-1 is purified by preparative HPLC and isolated as the trifluoroacetic acid salt. Mass spec.: M⁺ = 657/659.

EXAMPLE 25

20

Resin based synthesis of thrombin inhibitors

Step A: Preparation of Pro(p-nitrobenzophenoneoxime-polystyrene) resin

25 pNO₂ benzophenone-polystyrene oxime (0.5 mg/g, 1% cross-linked, 2.0 g) is slurried with BocProOH in 50 ml CH₂Cl₂ at room temperature and the suspension treated with 4 mL of a 0.5 M solution of dicyclohexylcarbodiimide in CH₂Cl₂. The mixture is shaken 24 hr at room temperature then filtered. The resin is washed with
30 alternating CH₂Cl₂ and ethylacetate and dried by suction.

The resin is suspended in a mixture of 15 ml trifluoroacetic acid and 30 ml of CH₂Cl₂ for 1.5 hr at room temperature and filtered. The resin is alternately steeped in CH₂Cl₂ and isopropanol then washed

- 40 -

with isopropanol and excess CH_2Cl_2 and dried to constant weight under vacuum; 2.0 g.

Step B: Preparation of Boc-D- β , β -diphenylala-Pro(p-nitrobenzophenoneoxime-polystyrene) resin

5 The resin from Step A is suspended in 20 ml CH_2Cl_2 containing 0.15 ml triethylamine and treated with a filtered solution of Boc D- β , β -diphenylalanine (1.02 g) in CH_2Cl_2 and 3 ml 0.5 M dicyclohexylcarbodiimide (removes dicyclohexylurea). The mixture is
10 shaken overnight at room temperature then filtered and washed alternating with isopropanol and CH_2Cl_2 and vacuum dried at 80°C. Amino acid analysis of the dried resin gave 214.8 mMol/g of Pro and an essentially equal amount of D- β , β -diphenylAla (after standard hydrolysis).

15

Step C: Release of dipeptide amides from resin and deblocking

 A 10 μMol equivalent of the resin from Step B is shaken with 2 ml CH_2Cl_2 containing an amine, preferably a benzylamine (10-13 μMol or its HCl salt and 100 μMol of triethylamine) for 24 hr at
20 room temperature. The mixture is filtered and the filtrate analyzed by HPLC to show the presence of Boc dipeptide amide and unreacted amine in constant ratio. The filtrates are concentrated under high vacuum and the residues treated with 10-20% trifluoroacetic acid in CH_2Cl_2 for 12
25 hr at room temperature. The mixtures are evaporated in a stream of nitrogen or under vacuum and the residues taken up in DMSO-water mixtures for bioassay as thrombin inhibitors.

- 41 -

EXAMPLE 26

Preparation of D-cyclohexylglycine-proline-N-(2-{O-carboxymethyl-N-ethylamide},5-chloro)-benzylamide (26-4)

5

Step A: Preparation of Boc-D-cyclohexylglycine-proline methyl ester (26-1)

A solution of 8.0 g (31.0 mmol) of Boc-D-cyclohexylglycine and 5.8 g (35 mmol) of proline methyl ester HCl salt in 100 ml of anh. DMF, mixed with 5.8 g (37.2 mmol) of HOBt with the pH adjusted to 7-8 with N-methylmorpholine (to moistened narrow-range pH paper), was treated with 7.9 g (40.3 mmol) of EDC and stirred for 18 hr in a nitrogen atmosphere. After 20 hr water (10 ml) was added, the solution concentrated *in vacuo* and partitioned with 400 ml EtOAc and 200 ml H₂O, washing with dil. NaHCO₃, H₂O, dil. KHSO₄, and twice with 50% satd NaCl, dried over Na₂SO₄ and concentrated under reduced pressure to give an oil. This crude material was chromatographed on 300 g silica gel in 1:1 (v/v) EtOAc/hexane to afford, after pooling of fractions, intermediary 26-1.

20

Step B: Preparation of Boc-D-cyclohexylglycine-proline (26-2)

26-1 (9.20 g) was dissolved in 90 ml of THF, adding 50 ml of H₂O, followed by 21 ml of 2.0 N LiOH in portions over a period of 2 hr. The solution was let stir 20 hr and the reaction was worked up by addition of dil. KHSO₄ to neutrality, evacuation of solvent under reduced pressure to give a thick paste to which was added 200 ml of H₂O in portions with stirring, followed by dil. KHSO₄ to acidity (pH < 2). After stirring for 1 hr, the solid was isolated by filtration, washing with H₂O twice, and drying *in vacuo* to give 6.45 g (72% yield overall) of intermediary Boc-D-cyclohexylglycine-proline. Evaporation of the filtrate to a volume of <100 ml afforded 26-2.

30

- 42 -

Step C: Preparation of Boc-D-cyclohexylglycine-proline-N-(2-
{O-carbethoxymethyl}-5-chloro)-benzylamide (26-3)

A solution of 405 mg (1.15 mmol) of 26-2 and 147 mg (0.94 mmol) of 2-hydroxy,5-chlorobenzylamine in 6 ml of anh. DMF, mixed with 191 mg (1.25 mmol) of HOBt with the pH adjusted to 7-8 with N-methylmorpholine (to moistened narrow-range pH paper), was treated with 255 mg (1.34 mmol) of EDC and stirred for 18 h in a nitrogen atmosphere. After 20 hr water (10 ml) was added, the solution concentrated *in vacuo* and partitioned with EtOAc and H₂O, washing with dil. NaHCO₃, H₂O, dil. KHSO₄, and twice with 50% satd NaCl, dried over Na₂SO₄ and concentrated under reduced pressure to give 502 mg of the crude 2-hydroxy,5-chlorobenzylamide.

A solution of this material in 20 ml of peroxide-free anh. dioxane was mixed with 0.18 ml (1.55 mmol) of ethyl bromoacetate and 0.54 g (1.66 mmol) of Cs₂CO₃ under a nitrogen atmosphere, stirring 20 hr at 25°. A second addition of 0.04 ml (0.34 mmol) of ethyl bromoacetate and 0.15 g (0.18 mmol) of Cs₂CO₃ brought the O-alkylation to completion, and the product was isolated by evaporation of solvent under reduced pressure, partitioning with EtOAc and H₂O, washing with dil. NaCl, drying over Na₂SO₄, and solvent removal to give 26-3.

Step D: Preparation of Boc-D-cyclohexylglycine-proline-N-(2-
{O-ethylacetamido}-5-chloro)-benzylamide (26-4)

26-3 (1.04g) was saponified in 30 ml of 50% THF/H₂O with 0.8 ml of 2.0 N LiOH for 20 hr, followed by addition of dil. KHSO₄ to neutrality, evaporation under reduced pressure to a gum, partitioning with EtOAc/dil. KHSO₄ and washing twice with dil. NaCl. After drying over Na₂SO₄, solvent removal afforded solid Boc-D-cyclohexylglycine-proline-N-(2-{O-carboxymethyl}-5-chloro)-benzylamide.

A solution of 91 mg (0.16 mmol) of the above acid and 35 mg (0.43 mmol) of ethylamine hydrochloride in 10 ml of anh. DMF, mixed with 37 mg (0.24 mmol) of HOBt with the pH adjusted to 7-8

- 43 -


with N-methylmorpholine (to moistened narrow-range pH paper), was treated with 58 mg (0.30 mmol) of EDC and stirred for 18 h in a nitrogen atmosphere. After 20 hr water (1 ml) was added, the solution concentrated *in vacuo* and partitioned with EtOAc and H₂O, washing
5 with dil. NaHCO₃, H₂O, dil. KHSO₄, and twice with 50% satd NaCl, dried over Na₂SO₄ and concentrated under reduced pressure to give the crude Boc-protected ethyl amide.

This intermediate was dissolved in 4 ml of 50% (v/v) TFA/CH₂Cl₂ for 30 min., the solvent was removed under reduced
10 pressure, and the product was purified by preparative HPLC (0.1% TFA-100% H₂O/CH₃CN -> 50% over 30 min.) to afford 26-4 as a white lyophilized powder. Anal. (C₂₄H₃₅N₄O₄Cl•1.30 TFA•0.15 H₂O), CHN. Mass spec.: M⁺ = 479.

The compounds shown in the tables below are exemplary
15 compounds of the present invention. The range of K_i values associated with the specifically listed compounds is represented as follows:

20
+++ <10 nM
++ >10 nM and <500 nM
+ >500 nM


- 44 -














TABLE I K_i

X	R	Thrombin	Trypsin
H		++	+
H		+++	+
H		++	+
H		++	+
H		++	+
$(\text{Me})_2\text{NCH}_2\text{CH}_2\text{OC}(=\text{O})-$		++	+
$-\text{CH}_2\text{COOH}$		++	+


- 45 -

TABLE I CONT'D




X	R	K _i	
		Thrombin	Trypsin
		++	+
		++	+
		++	+
		++	+
HOCH ₂ CH ₂ -		++	+
		+++	+
-CH ₂ CON(Et) ₂		++	+
-CH ₂ CON(Et) ₂		+++	+

- 46 -

TABLE 1 CONT'D


X	R	K _i	
		Thrombin	Trypsin
		+++	+
		+++	+
		+++	+
		+++	+
 -CH ₂ COOH		++	+

- 47 -

TABLE II


R	Thrombin	K_i	Trypsin
<p>$\text{H}_2\text{N}-\text{CH}(\text{CO})-\text{CH}_2-\text{C}_6\text{H}_3\text{O}_2$</p>	+++		+
<p>$\text{H}_2\text{N}-\text{CH}(\text{CO})-\text{CH}(\text{OH})-\text{C}_6\text{H}_5$</p>	++		+
<p>$\text{H}_2\text{N}-\text{CH}(\text{CO})-\text{CH}_2-\text{C}_6\text{H}_4\text{OMe}$</p>	++		+
<p>$\text{H}_2\text{N}-\text{CH}(\text{CO})-\text{CH}_2-\text{C}_6\text{H}_5$</p>	++		+

- 48 -

TABLE III


R	K _i	
	Thrombin	Trypsin
H	+++	+
CH ₂ COOH	+++	+
CH ₂ CON(Et) ₂	+++	+
	++	+
	+++	+




- 49 -

TABLE IV

R	Thrombin	Trypsin
-OEt	+++	+
-OH	+++	+
-NH-Et	+++	+
-NH ₂	+++	+


- 50 -


TABLE IV (CON'T)

R	Thrombin	Trypsin
-NH (CH ₂) ₂ OH	+++	+
-NH- 	+++	+
-N-  -OH	+++	+
-N- 	++	+

- 51 -

TABLE V



R _a	R _b	D	K _i	
			Thrombin	Trypsin
CH ₃	H	H	++	+
CH ₃	CH ₃	H	++	+
H	CH ₂ CH ₃	H	++	+
H	CH ₂ CH ₂ CH ₃	H	+++	+
CH ₂ CH ₃	CH ₂ CH ₃	H	+++	+
CH ₂ CH ₃	CH ₂ CH ₃	CH ₃	+++	+
CH ₂ CH ₂ CH ₃	CH ₂ CH ₂ CH ₃	H	+++	+
H	CH ₂ CH(CH ₃) ₂	H	+++	+
H	CH ₂ CH 	H	+++	+

- 52 -

K_i for thrombin range is the inhibition constant for the tested compound with human thrombin. K_i for trypsin is the inhibition constant for the tested compound with human trypsin. Rate constants were determined using the following *in vitro* procedures.

5

In vitro assay for determining proteinase inhibition

Assays of human α -thrombin and human trypsin were performed at 25°C in 0.05 M TRIS buffer pH 7.4, 0.15 M NaCl, 0.1% PEG. Trypsin assays also contained 1 mM CaCl_2 .

In assays wherein rates of hydrolysis of a *p*-nitroanilide (pna) substrate were determined, a Thermomax 96-well plate reader was used to measure (at 405 nm) the time dependent appearance of *p*-nitroaniline. sar-PR-pna (sarcosine-Pro-Arg-*p*-nitroanilide) was used to assay human α -thrombin ($K_M=125 \mu\text{M}$) and human trypsin ($K_M=59 \mu\text{M}$). *p*-Nitroanilide substrate concentration was determined from measurements of absorbance at 342 nm using an extinction coefficient of $8270 \text{ cm}^{-1}\text{M}^{-1}$.

In certain studies with potent inhibitors ($K_i < 10 \text{ nM}$) where the degree of inhibition of thrombin was high, a more sensitive activity assay was employed. In this assay the rate of thrombin catalyzed hydrolysis of the fluorogenic substrate Z-GPR-afc (Cbz-Gly-Pro-Arg-7-amino-4-trifluoromethyl coumarin) ($K_M=27 \mu\text{M}$) was determined from the increase in fluorescence at 500 nm (excitation at 400 nm) associated with production of 7-amino-4-trifluoromethyl coumarin. Concentrations of stock solutions of Z-GPR-afc were determined from measurements of absorbance at 380 nm of the 7-amino-4-trifluoromethyl coumarin produced upon complete hydrolysis of an aliquot of the stock solution by thrombin.

Activity assays were performed by diluting a stock solution of substrate at least tenfold to a final concentration $\leq 0.5 K_M$ into a solution containing enzyme or enzyme equilibrated with inhibitor. Times required to achieve equilibration between enzyme and inhibitor were determined in control experiments. Initial velocities of product

- 53 -

formation in the absence (V_o) or presence of inhibitor (V_i) were measured. Assuming competitive inhibition, and that unity is negligible compared $K_m/[S]$, $[I]/e$, and $[I]/e$ (where $[S]$, $[I]$, and e respectively represent the total concentrations, of substrate, inhibitor and enzyme), the equilibrium constant (K_i) for dissociation of the inhibitor from the enzyme can be obtained from the dependence of V_o/V_i on $[I]$ shown in equation 1.

$$V_o/V_i = 1 + [I]/K_i \quad (1)$$

10

The activities shown by this assay indicate that the compounds of the invention are therapeutically useful for treating various conditions in patients suffering from unstable angina, refractory angina, myocardial infarction, transient ischemic attacks, atrial fibrillation, thrombotic stroke, embolic stroke, deep vein thrombosis, disseminated intravascular coagulation, and reocclusion or restenosis of recanalized vessels.

15

Thrombin Inhibitors - Therapeutic Uses

20

Anticoagulant therapy is indicated for the treatment and prevention of a variety of thrombotic conditions, particularly coronary artery and cerebrovascular disease. Those experienced in this field are readily aware of the circumstances requiring anticoagulant therapy. The term "patient" used herein is taken to mean mammals such as primates, including humans, sheep, horses, cattle, pigs, dogs, cats, rats, and mice.

25

Thrombin inhibition is useful not only in the anticoagulant therapy of individuals having thrombotic conditions, but is useful whenever inhibition of blood coagulation is required such as to prevent coagulation of stored whole blood and to prevent coagulation in other biological samples for testing or storage. Thus, thrombin inhibitors can be added to or contacted with any medium containing or suspected of containing thrombin and in which it is desired that blood coagulation be inhibited, e.g. when contacting the mammal's blood with material

30

- 54 -

selected from the group consisting of vascular grafts, stents, orthopedic prosthesis, cardiac prosthesis, and extracorporeal circulation systems

The thrombin inhibitors of the invention can be administered in such oral forms as tablets, capsules (each of which
5 includes sustained release or timed release formulations), pills, powders, granules, elixers, tinctures, suspensions, syrups, and emulsions. Likewise, they may be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using forms well known to those of ordinary skill in the pharmaceutical arts. An
10 effective but non-toxic amount of the compound desired can be employed as an anti-aggregation agent. For treating ocular build up of fibrin, the compounds may be administered intraocularly or topically as well as orally or parenterally.

The thrombin inhibitors can be administered in the form of
15 a depot injection or implant preparation which may be formulated in such a manner as to permit a sustained release of the active ingredient. The active ingredient can be compressed into pellets or small cylinders and implanted subcutaneously or intramuscularly as depot injections or implants. Implants may employ inert materials such as biodegradable
20 polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers manufactured by the Dow-Corning Corporation.

The thrombin inhibitors can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be
25 formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

The thrombin inhibitors may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The thrombin inhibitors may also be coupled
30 with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the thrombin inhibitors may be coupled to a class of

- 55 -

biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihdropyrans, 5 polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels.

The dosage regimen utilizing the thrombin inhibitors is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the 10 condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the condition.

15 Oral dosages of the thrombin inhibitors, when used for the indicated effects, will range between about 0.1 mg per kg of body weight per day (mg/kg/day) to about 100 mg/kg/day and preferably 1.0-100 mg/kg/day and most preferably 1-20 mg/kg/day. Intravenously, the most preferred doses will range from about 0.01 to about 10 20 mg/kg/minute during a constant rate infusion. Advantageously, the thrombin inhibitors may be administered in divided doses of two, three, or four times daily. Furthermore, they can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well 25 known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, or course, be continuous rather than intermittent throughout the dosage regime.

For example, oral tablets can be prepared which contain an 30 amount of active compound of between 25 and 500 mg, typically between 200 and 250 mg. Typically, a patient in need of thrombin inhibitor compound, depending on weight and metabolism of the patient, would be administered between about 100 and 1000 mg active compound per day. For a patient requiring 1000 mg per day, two

- 56 -

tablets containing 250 mg of active compound can be administered in the morning and two tablets containing 250 mg of active compound can again be administered in the evening. For a patient requiring 500 mg per day, one tablet containing 250 mg of active compound can be
5 administered in the morning and one tablet containing 250 mg of active compound can again be administered in the evening.

The thrombin inhibitors are typically administered as active ingredients in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as "carrier"
10 materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixers, syrups and the like, and consistent with convention pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral,
15 non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert
20 carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn-sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene
25 glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch methyl cellulose, agar, bentonite,
30 xanthan gum and the like.

The thrombin inhibitors can also be co-administered with suitable anti-coagulation agents or thrombolytic agents such as plasminogen activators or streptokinase to achieve synergistic effects in the treatment of various ascular pathologies. For example, thrombin

- 57 -

inhibitors enhance the efficiency of tissue plasminogen activator-mediated thrombolytic reperfusion. Thrombin inhibitors may be administered first following thrombus formation, and tissue plasminogen activator or other plasminogen activator is administered thereafter. They may also be combined with heparin, aspirin, or warfarin.

EXAMPLE 27

Tablet Preparation

Tablets containing 25.0, 50.0, and 100.0 mg, respectively, of the following active compounds are prepared as illustrated below:

N-[4-(imidazolyl)-methyl]-D- β , β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide

N-[2-(5-hydroxymethylfuryl)-methyl]-D- β , β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide

N-[2-(5-dimethylaminofuryl)-methyl]-D- β , β -diphenylala-Pro-N-(2,5-dichloro)-benzylamide

	<u>Ingredient</u>	<u>Amount-mg</u>		
		25.0	50.0	100.0
25	Microcrystalline cellulose	37.25	100.0	200.0
	Modified food corn starch	37.25	4.25	8.5
30	Magnesium stearate	0.50	0.75	1.5

All of the active compound, cellulose, and a portion of the corn starch are mixed and granulated to 10% corn starch paste. The resulting granulation is sieved, dried and blended with the remainder of the corn starch and the magnesium stearate. The resulting granulation is

- 58 -

then compressed into tablets containing 25.0, 50.0, and 100.0 mg, respectively, of active ingredient per tablet.

EXAMPLE 28

5

An intravenous dosage form of the above-indicated active compound is prepared as follows:

10	Active Compound	0.5-10.0mg
	Sodium Citrate	5-50mg
	Citric Acid	1-15mg
15	Sodium Chloride	1-8mg
	Water for Injection (USP)	q.s. to 1 L


20 Utilizing the above quantities, the active compound is dissolved at room temperature in a previously prepared solution of sodium chloride, citric acid, and sodium citrate in Water for Injection (USP, see page 1636 of United States Pharmacopeia/National Formulary for 1995, published by United States Pharmacopeial Convention, Inc., Rockville, Maryland, copyright 1994.

25

- 59 -

WHAT IS CLAIMED IS:

1. A compound having the following structure:




5

I

and pharmaceutically acceptable salts thereof wherein

A is

10



wherein

15

R_a and R_b are independently selected from
hydrogen,

20

a heterocyclic group which is a stable 5- to 7-membered
mono- or bicyclic or stable 7- to 10-membered bicyclic
heterocyclic ring system any ring of which may be
saturated or unsaturated, and which consists of carbon
atoms and from one to three heteroatoms selected from the
group consisting of N, O and S, and wherein the nitrogen
and sulfur heteroatoms may optionally be oxidized, and the
nitrogen heteroatom may optionally be quaternized, and
including any bicyclic group in which any of the above-
defined heterocyclic rings is fused to a benzene ring,

25

- 60 -

C₁₋₄ alkyl unsubstituted or substituted with CH₃ or C₃₋₇ cycloalkyl,

aryl,

substituted aryl with one or two substituents selected from

5

C₁₋₄ alkyl,

C₁₋₄ alkoxy,

methylenedioxy,

halogen or


hydroxy,

10

C₃₋₇ cycloalkyl,

a C₄₋₁₀ carbocyclic or bicyclic ring, or

R_a and R_b, along with the carbon to which they are attached, form a C₃₋₇ cycloalkyl ring or



15

where R¹⁰ is H or -OH, and

R¹¹ is H or -OCH₃, and

X is -NHR_C or -OH, wherein,

20

R_C is

hydrogen,

-CH₃,

-(CH₂)₁₋₃CH₃,

-(CH₂)₂₋₄OH,

25


-(CH₂)₁₋₃COOH,

-(CH₂)₁₋₃COOR⁶, where R⁶ is C₁₋₄alkyl,

-(CH₂)₁₋₃CONR⁷R⁸,

where R⁷ and R⁸ are independently hydrogen or C₁₋₄alkyl,

30




- 61 -

where D is 1, 2, 3, or 4 carbon atoms unsubstituted
or any 1, 2, 3, or 4 of which are substituted with OH,
-SO₂(CH₂)₁₋₃aryl,
-(CH₂)₁₋₃NH₂,

5

C₃₋₇ cycloalkyl ring unsubstituted or substituted with
-OH, -C(O)OH, or -C(O)OR_d, where R_d is
C₁₋₄ alkyl,




10

Y is O or NH,

W is C or N,

Z is C or N, and


R⁶ is -CH₂OH or -N(CH₃)₂ provided that W and Z
are not the same,






15

R⁷ is H or CH₃, and


R⁸ is H or



20




- 62 -



where R^9 is H, NH_2 , or OH;

5 or

A is



, wherein

10

B is a bond, O, $-\text{CH}_2-\text{O}-$, or $-\text{O}-\text{CH}_2-$;

R^2 and R^5 are independently selected from

hydrogen, provided that R^2 and R^5 are not both hydrogen,

15 C_{1-4} alkyl,

C_{1-4} alkoxy,

halogen,

$-\text{COOH}$,

$-\text{OH}$,

20 $-\text{COOR}^6$, where R^6 is C_{1-4} alkyl,

$-\text{CONR}^7\text{R}^8$, where R^7 and R^8 are independently
hydrogen or C_{1-4} alkyl,

$-\text{OCH}_2\text{CO}_2\text{H}$,

$-\text{OCH}_2\text{CO}_2\text{CH}_3$,

25 $-\text{OCH}_2\text{CO}_2(\text{CH}_2)_{1-3}\text{CH}_3$,

$-\text{O}(\text{CH}_2)_{1-3}\text{C}(\text{O})\text{NR}^3\text{R}^4$, wherein R^3 and R^4 are independently
hydrogen, C_{1-4} alkyl, C_{3-7} cycloalkyl, or $-\text{CH}_2\text{CF}_3$,

$-(\text{CH}_2)_{1-4}\text{OH}$,


$-\text{NHC}(\text{O})\text{CH}_3$.

- 63 -

-NHC(O)CF₃,
 -NHSO₂CH₃, and
 -SO₂NH₂; and

5 m is 1 or 2.

2. A compound of claim 1 having the following structure:




10

and pharmaceutically acceptable salts thereof.

3. A compound of claim 2 having the following structure:

15






and pharmaceutically acceptable salts thereof, wherein

20 R² is -OCH₂C(O)NHR⁴; and
 R⁴ is -CH₂CH₃, cyclopropyl, or -CH₂CF₃.

4. A compound of claim 3 selected from the group consisting of:

- 64 -




5

and pharmaceutically acceptable salts thereof.

5. A compound of claim 2 having the following
10 structure:

- 65 -



and pharmaceutically acceptable salts thereof wherein

5 X is -NHR_C or -OH, wherein

R_C is

hydrogen,

-CH₃,

10 -(CH₂)₁₋₃CH₃,


-(CH₂)₂₋₄OH,

-(CH₂)₁₋₃COOH,

-(CH₂)₁₋₃COOR⁶, where R⁶ is C₁₋₄alkyl,

-(CH₂)₁₋₃CONR⁷R⁸, where R⁷ and R⁸ are independently

15 hydrogen or C₁₋₄alkyl,



where D is 1, 2, 3, or 4 carbon atoms unsubstituted or


any 1, 2, 3, or 4 of which are substituted with OH,

-SO₂(CH₂)₁₋₃aryl,

20 -(CH₂)₁₋₃NH₂,

C₃₋₇ cycloalkyl ring unsubstituted or substituted with -OH,

-C(O)OH, or -C(O)OR_d, where R_d is C₁₋₄ alkyl,



where


25

Y is O or NH,

- 66 -


W is C or N,

Z is C or N, and




R⁶ is -CH₂OH or -N(CH₃)₂ provided that W and Z are not the same,

5

where

R⁷ is H or CH₃, andR⁸ is H or

10



15

where R⁹ is H, NH₂, or OH;R² and R⁵ are independently selected fromhydrogen, provided that R² and R⁵ are not both hydrogen,C₁₋₄ alkyl,

20


C₁₋₄ alkoxy,

halogen, and

-OH.

- 67 -

6. A compound of claim 5 having the following structure:



5

and pharmaceutically acceptable salts thereof wherein

R_a and R_b are independently selected from

hydrogen,

10

a heterocyclic group which is a stable 5- to 7-membered mono- or bicyclic or stable 7- to 10-membered bicyclic heterocyclic ring system any ring of which may be saturated or unsaturated, and which consists of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O and S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring,

15

20

C₁₋₄ alkyl unsubstituted or substituted with CH₃ or C₃₋₇ cycloalkyl, phenyl, or

R_a and R_b , along with the carbon to which they are attached, form a cyclohexyl ring; and

25


R^2 and R^5 are independently selected from hydrogen, provided that R^2 and R^5 are not both hydrogen, Cl,

- 68 -

-CH₃,
 -CH₂CH₃,
 -OCH₃, and
 -OH.

5

7. A compound of claim 6 having the following structure:




10

and pharmaceutically acceptable salts thereof wherein

R² and R⁵ are independently selected from -OCH₃ and -CH₃; and



15 R_C is hydrogen or -SO₂CH₂C₆H₅.

8. A compound of claim 7 selected from the group consisting of:



20


- 69 -



and pharmaceutically acceptable salts thereof.

5

9. A compound of claim 6 having the following structure:




10

and pharmaceutically acceptable salts thereof.

10. A compound of claim 9 having the following structure:

15

- 70 -




and pharmaceutically acceptable salts thereof, wherein

5 R_c is

hydrogen,

$SO_2CH_2C_6H_5$, or

10






; and

R_a and R_b are phenyl, or R_a and R_b , along with the carbon to which they are attached, form cyclohexyl.



15

11. A compound of claim 10 selected from the group consisting of:

- 71 -



- 72 -



and pharmaceutically acceptable salts thereof.

5

12. A composition for inhibiting thrombin in blood comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

10

13. A composition for inhibiting thrombus formation in blood comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

15

14. A method for inhibiting thrombin in blood in a mammal comprising administering to the mammal a composition of Claim 12.

20

15. A method for inhibiting thrombus formation in blood in a mammal comprising administering to the mammal a composition of Claim 13.

- 73 -

16. A method for inhibiting thrombin in stored blood comprising administering to the mammal a composition of Claim 12.

5 17. A method for inhibiting thrombus formation in stored blood comprising administering to the mammal a composition of Claim 13.

10 18. A composition for inhibiting thrombus formation in blood comprising a compound of Claim 1, a fibrinogen receptor antagonist, and a pharmaceutically acceptable carrier.

15 19. A method for inhibiting thrombus formation in blood in a mammal comprising administering to the mammal a composition of Claim 18.

20 20. The use of a compound of Claim 1, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for inhibiting thrombus formation, preventing thrombus formation, inhibiting thrombin, inhibiting formation of fibrin, and inhibiting formation of blood platelet aggregates, in a mammal.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/16865

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :A01N 43/34, 43/64, 43/82; A61K 31/165; C07D 207/08 US CL :514/359; 548/566 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/359; 548/566 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EDWARDS et al. Design, Synthesis, and Kinetic Evaluation of a Unique Class of Elastase Inhibitors, the Peptidyl α -Ketobenzoxazoles, and the X-ray Crystal Structure of the Covalent Complex between Porcine Pancreatic Elastase and Ac-Ala-Pro-Val-2-Benzoxazole. J. Am. Chem. Soc. 26 February 1992, Vol. 114, No. 5, pages 1854-1863.	1-11
A	EP 0 363 284 A2 (MERRELL DOW PHARMACEUTICALS INC.) 11 April 1990 (11.04.90), see pages 2-3.	1-20
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search 04 DECEMBER 1996		Date of mailing of the international search report 21 JAN 1997
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer JANE OSWECKI <i>Jab for</i> Telephone No. (703) 308-1235